



Parent/Athlete Concussion Information Sheet

A concussion is a type of traumatic brain injury that changes the way the brain normally works. A concussion is caused by bump, blow, or jolt to the head or body that causes the head and brain to move rapidly back and forth. Even a “ding,” “getting your bell rung,” or what seems to be a mild bump or blow to the head can be serious.

WHAT ARE THE SIGNS AND SYMPTOMS OF CONCUSSION?

Signs and symptoms of concussion can show up right after the injury or may not appear or be noticed until days or weeks after the injury.

If an athlete reports **one or more** symptoms of concussion listed below after a bump, blow, or jolt to

Did You Know?

- Most concussions occur *without* loss of consciousness.
- Athletes who have, at any point in their lives, had a concussion have an increased risk for another concussion.
- Young children and teens are more likely to get a concussion and take longer to recover than adults.

the head or body, s/he should be kept out of play the day of the injury and until a health care professional, experienced in evaluating for concussion, says s/he is symptom-free and it’s OK to return to play.

SIGNS OBSERVED BY COACHING STAFF	SYMPTOMS REPORTED BY ATHLETES
Appears dazed or stunned	Headache or “pressure” in head
Is confused about assignment or position	Nausea or vomiting
Forgets an instruction	Balance problems or dizziness
Is unsure of game, score, or opponent	Double or blurry vision
Moves clumsily	Sensitivity to light
Answers questions slowly	Sensitivity to noise
Loses consciousness (<i>even briefly</i>)	Feeling sluggish, hazy, foggy, or groggy
Shows mood, behavior, or personality changes	Concentration or memory problems
Can’t recall events <i>prior</i> to hit or fall	Confusion
Can’t recall events <i>after</i> hit or fall	Just not “feeling right” or “feeling down”

CONCUSSION DANGER SIGNS

In rare cases, a dangerous blood clot may form on the brain in a person with a concussion and crowd the brain against the skull. An athlete should receive immediate medical attention if after a bump, blow, or jolt to the head or body s/he exhibits any of the following danger signs:

- One pupil larger than the other
- Is drowsy or cannot be awakened
- A headache that not only does not diminish, but gets worse
- Weakness, numbness, or decreased coordination
- Repeated vomiting or nausea
- Slurred speech
- Convulsions or seizures
- Cannot recognize people or places
- Becomes increasingly confused, restless, or agitated
- Has unusual behavior
- Loses consciousness (*even a brief loss of consciousness should be taken seriously*)

WHY SHOULD AN ATHLETE REPORT THEIR SYMPTOMS?

If an athlete has a concussion, his/her brain needs time to heal. While an athlete's brain is still healing, s/he is much more likely to have another concussion. Repeat concussions can increase the time it takes to recover. In rare cases, repeat concussions in young athletes can result in brain swelling or permanent damage to their brain. *They can even be fatal.*

Remember

Concussions affect people differently. While most athletes with a concussion recover quickly and fully, some will have symptoms that last for days, or even weeks. A more serious concussion can last for months or longer.

WHAT SHOULD YOU DO IF YOU THINK YOUR ATHLETE HAS A CONCUSSION?

If you suspect that an athlete has a concussion, remove the athlete from play and seek medical attention. Do not try to judge the severity of the injury yourself. Keep the athlete out of play the day of the injury and until a health care professional, experienced in evaluating for concussion, says s/he is symptom-free and it's OK to return to play.

Rest is key to helping an athlete recover from a concussion. Exercising or activities that involve a lot of concentration, such as studying, working on the computer, or playing video games, may cause concussion symptoms to reappear or get worse. After a concussion, returning to sports and school is a gradual process that should be carefully managed and monitored by a health care professional.

It's better to miss one game than the whole season. For more information on concussions, visit: www.cdc.gov/Concussion.



SPORT SCIENCE
INSTITUTE™



CONCUSSION SAFETY

WHAT STUDENT-ATHLETES
NEED TO KNOW

What is a concussion?

A concussion is a type of traumatic brain injury. It follows a force to the head or body and leads to a change in brain function. It is not typically accompanied by loss of consciousness.

How can I keep myself safe?

1. Know the symptoms.

You may experience ...

- Headache or head pressure
- Nausea
- Balance problems or dizziness
- Double or blurry vision
- Sensitivity to light or noise
- Feeling sluggish, hazy or foggy
- Confusion, concentration or memory problems

2. Speak up.

- If you think you have a concussion, stop playing and talk to your coach, athletic trainer or team physician immediately.

3. Take time to recover.

- Follow your team physician and athletic trainer's directions during concussion recovery. If left unmanaged, there may be serious consequences.
- Once you've recovered from a concussion, talk with your physician about the risks and benefits of continuing to participate in your sport.

How can I be a good teammate?

1. Know the symptoms.

You may notice that a teammate ...

- Appears dazed or stunned
- Forgets an instruction
- Is confused about an assignment or position
- Is unsure of the game, score or opponent
- Appears less coordinated
- Answers questions slowly
- Loses consciousness

2. Encourage teammates to be safe.

- If you think one of your teammates has a concussion, tell your coach, athletic trainer or team physician immediately.
- Help create a culture of safety by encouraging your teammates to report any concussion symptoms.

3. Support your injured teammates.

- If one of your teammates has a concussion, let him or her know you and the team support playing it safe and following medical advice during recovery.
- Being unable to practice or join team activities can be isolating. Make sure your teammates know they're not alone.

No two concussions are the same. New symptoms can appear hours or days after the initial impact. If you are unsure if you have a concussion, talk to your athletic trainer or team physician immediately.

What happens if I get a concussion and keep practicing or competing?

- Due to brain vulnerability after a concussion, an athlete may be more likely to suffer another concussion while symptomatic from the first one.
- In rare cases, repeat head trauma can result in brain swelling, permanent brain damage or even death.
- Continuing to play after a concussion increases the chance of sustaining other injuries too, not just concussion.
- Athletes with concussion have reduced concentration and slowed reaction time. This means that you won't be performing at your best.
- Athletes who delay reporting concussion take longer to recover fully.

What are the long-term effects of a concussion?

- We don't fully understand the long-term effects of a concussion, but ongoing studies raise concerns.
- Athletes who have had multiple concussions *may* have an increased risk of degenerative brain disease and cognitive and emotional difficulties later in life.

What do I need to know about repetitive head impacts?

- Repetitive head impacts mean that an individual has been exposed to repeated impact forces to the head. These forces may or may not meet the threshold of a concussion.
- Research is ongoing but emerging data suggest that repetitive head impact also may be harmful and place a student-athlete at an increased risk of neurological complications later in life.

Did you know?

- NCAA rules require that team physicians and athletic trainers manage your concussion and injury recovery independent of coaching staff, or other non-medical, influence.
- We're learning more about concussion every day. To find out more about the largest concussion study ever conducted, which is being led by the NCAA and U.S. Department of Defense, visit ncaa.org/concussion.

CONCUSSION TIMELINE

