



Athletes who experience one or more of the signs and symptoms listed below after a bump, blow, or jolt to the head or body may have a concussion.

## SYMPTOMS REPORTED BY ATHLETE:

- Headache or "pressure" in head
- Nausea or vomiting
- Balance problems or dizziness
- Double or blurry vision
- Sensitivity to light
- Sensitivity to noise
- Feeling sluggish, hazy, foggy, or groggy
- Concentration or memory problems
- Confusion
- Just not "feeling right" or is "feeling down"

## SIGNS OBSERVED BY COACHING STAFF:

- Appears dazed or stunned
- Is confused about assignment or position
- Forgets an instruction
- Is unsure of game, score, or opponent
- Moves clumsily
- Answers questions slowly
- Loses consciousness (even briefly)
- Shows mood, behavior, or personality changes
- Can't recall events prior to hit or fall
- Can't recall events after hit or fall

## IF YOU SUSPECT THAT AN ATHLETE HAS A CONCUSSION, YOU SHOULD TAKE THE FOLLOWING STEPS:

### HEADS UP ACTION PLAN:

1. Remove the athlete from play.
2. Keep the athlete out of play the day of the injury.
3. Obtain permission from an appropriate health care professional that states the athlete can return to play.

## IMPORTANT PHONE NUMBERS:

### EMERGENCY MEDICAL SERVICES

NAME: \_\_\_\_\_

PHONE: \_\_\_\_\_

### HEALTH CARE PROFESSIONAL

NAME: \_\_\_\_\_

PHONE: \_\_\_\_\_

### STAFF AVAILABLE DURING PRACTICES

NAME: \_\_\_\_\_

PHONE: \_\_\_\_\_

### STAFF AVAILABLE DURING GAMES

NAME: \_\_\_\_\_

PHONE: \_\_\_\_\_